

## REVIEW ARTICLE

# Modulation of microRNAs in Tooth Root and Periodontal Tissue Development

Ran Zhang\* and Tiejun Li

Department of Oral Pathology, School and Hospital of Stomatology, Peking University, Beijing, China

## ARTICLE HISTORY

Received: December 06, 2016  
Revised: June 14, 2017  
Accepted: July 01, 2017

DOI:  
10.2174/1574888X12666170713123647

**Abstract:** Tooth root development begins after the completion of tooth crown development. Both the tooth root and crown undergo a series of interactions between the epithelium and adjacent mesenchymal cells. Although many studies have evaluated tooth crown formation, little is known about the regulatory mechanisms of tooth root development. MicroRNAs (miRNAs) are small noncoding RNAs that regulate protein expression through post-transcriptional mechanisms and participate in a broad range of biological processes, from development to tumorigenesis. In this review, we summarize current knowledge on the role of miRNAs in regulating the morphogenesis of root odontoblasts and periodontal tissues and regarding the interplay between miRNAs and signaling molecules during these processes.

**Keywords:** microRNA, root odontogenesis, periodontal tissue.

## 1. INTRODUCTION

Tooth regeneration is a promising approach based on a comprehensive understanding of reciprocal interactions between the epithelium and underlying mesenchymal cells [1]. Unlike crown development, tooth root development begins postnatally and is a consequence of sequential crosstalk between Hertwig's epithelial root sheath and mesenchymal cells. Dysregulation of these interactions leads to malformed tooth roots or abnormal periodontal structures [2]. A variety of mesenchymal molecules contribute to root odontoblast and cementoblast differentiation and periodontal tissue construction during tooth root development [3-7]. Previous studies have implied that signals from members of the Wnt family, transforming growth factor  $\beta$  (TGF- $\beta$ ) superfamily, and sonic hedgehog (SHH) pathway are involved in the epithelial-mesenchymal interactions required for regulating tooth root development [4, 7-9]. Transcription factors such as nuclear factor 1 C (NF1C), osterix, and runt-related transcription factor 2 (RUNX2) govern the expression of proteins in these signaling pathways and play important roles during tooth root development [4, 10-12].

MicroRNAs (miRNAs) are small, noncoding, single-stranded RNAs with approximately 18-22 nucleotides that regulate gene expression post-transcriptionally. miRNAs are transcribed from endogenous miRNA genes and generate primary (pri-) miRNAs [13, 14]. Pri-miRNAs are processed into single hairpins or precursor miRNAs (pre-miRNAs) by

the RNAase III enzyme in the nucleus. The pre-miRNAs undergo cleavage by a second RNA III enzyme, Dicer, after translocation of pre-miRNAs into the cytoplasm. Eventually, the mature miRNAs function in the form of an RNA-induced silencing complex and regulate the expression of multiple targets by binding to the 3'untranslated region (UTR) [15, 16]. One miRNA can target several genes, and several miRNAs can target a single gene [17, 18]. Thus, miRNAs act as potent molecular managers that can control multiple developmental and physiological processes [19, 20].

miRNAs plays an essential role in both bone and tooth formation and are upstream regulators of several signaling pathways and transcription factors, such as RUNX2 and osterix [21]. Recently, the important and unique role of miRNAs in tooth root formation has been gradually realized [22]. In this review, we summarize recent discoveries of the biological functions of miRNAs in root odontogenesis, the development of periodontal tissues and stem cells from dental or periodontal tissues, providing a better understanding of clinical tooth root malformation and potential therapeutic approaches.

## 2. TOOTH ROOT AND PERIODONTAL TISSUE DEVELOPMENT

Tooth crown development begins in the late embryonic period with epithelial invaginations from the alveolar mucosa, also known as the dental lamina. The dental lamina gives rise to enamel organs and subsequently induced the underlying mesenchymal cells to differentiate and form the structure of dental papilla and dental sac [1]. Crown development is completed through the interaction between the

\*Address correspondence to this author at the Department of Oral Pathology, School and Hospital of Stomatology, Peking University, Beijing, China; E-mail: 15404026@qq.com

dental lamina and dental papilla, which occur via amelogenesis and crown odontogenesis [23]. With the establishment of conditional knockout mice, it is possible to manipulate specific genes in the dental epithelium or mesenchyme and observe *in vivo* functions of numerous genes in tooth morphogenesis. Members of various signaling pathways, such as the TGF- $\beta$  superfamily, Wnt, Notch, and Shh, have indispensable roles in amelogenesis and crown odontogenesis [24-27].

The tooth root development process begins with the extension of Hertwig's epithelial root sheath (HERS), a bilayered structure formed by inner and outer enamel epithelium. HERS acts as an inducer, expressing signaling molecules to promote differentiation of apical papilla into root odontoblasts, which later will produce root dentin. With the formation of root dentin, HERS breaks into nests and cords, leaving space for the outer mesenchymal cells to contact the newly formed root dentin and posteriorly differentiate into cementoblasts and periodontal tissues [28] (Fig. 1). The crown is formed during early embryonic development, whereas the root is formed from the late embryonic stage to the postnatal stage (up to postnatal day 30.5 in mice) [29]. Few animal models can survive after birth, particularly those with deficiencies in critical genes; thus, tooth root development has not been sufficiently studied.


### 3. FUNCTION OF MIRNAS IN TOOTH ROOT AND PERIODONTAL TISSUE FORMATION

#### 3.1. Function of miRNAs in Root Odontoblasts

Crown and root odontogenesis were previously thought to undergo similar mechanisms. However, in NF1C-null mice, root dentin development was blocked, while crown dentin formation was unaffected [30]. In the recent years, with the emergence of *Cre* transgenic lines that effectively ablate genes within tooth root odontoblasts or HERS, signaling pathways involved in manipulating the formation and

development of root dentin have been elucidated [7, 31, 32]. Signaling molecules of the TGF- $\beta$  superfamily and Wnt family as well as transcription factors such as RUNX2, osterix, and NF1C are also involved in root odontoblast differentiation. Thus, miRNAs targeting these proteins may be vital for dentin formation.

Bone morphogenic protein (BMP) signaling, from both the apical papilla and HERS, has been shown to be critical in tooth root development [3, 31]. Moreover, miRNAs have been reported to regulate odontogenesis by modulating the BMP signaling pathway [33, 34]. In cultured human dental papilla cells, BMP7 protein is downregulated after transfection with an miR-34a mimic, but was upregulated after transfection with an miR-34a inhibitor. A previous study concluded that miR-34 may inhibit the expression of alkaline phosphatase (ALP) and upregulate dentine sialophosphoprotein (DSPP) by targeting BMP7 [35]. A recent study also showed that miR-135b binds to the 3'-UTR of *Smad5* and *Smad4* mRNAs and suppresses their expression, consequently blocking the differentiation of human dental pulp cells [36]. Our previous data suggested that Wnt/ $\beta$ -catenin signaling is functionally significant for root odontogenesis [7]. Notably, both miR-663 and miR-27 have binding sites in the 3' UTR of the adenomatous polyposis coli (APC) gene. Binding of either miR-663 or miR-27 with APC activates Wnt/ $\beta$ -catenin signaling in MDP-23 cells, promoting the differentiation of odontoblasts [37, 38]. RUNX2 is of vital importance in odontoblast differentiation, it shows higher expression in differentiating odontoblasts and decreased expression in differentiated odontoblast [39-41]. In RUNX2 deficient mice, the peripheral layer of cells in the dental papilla lost polarization and odontoblastic differentiation was blocked [42]. By overexpression Runx2 under the control of *Colla1* promoter, the terminal differentiation of odontoblasts was severely disturbed by showing flat or cuboid shape with reduced differentiation markers (*Colla1*, *DSPP*, nestin) [43]. According to a previous study, overexpression of miR-338-



**Fig. (1).** The illustration of tooth root development. (A) Initial stage. The inner and outer epitheliums constitute a structure named cervical loop, and began to extend into the apical direction. (B) Intermediate stage. The extension of HERS induces the odontoblastic differentiation of apical papilla cells. (C) Late stage. HERS cells break into nests and cords allowing the dental follicle cells to contact the newly-formed dentin and then form the periodontal structures.

3p decreases the expression of RUNX2, while miR-338-3p inhibitors increase the expression of RUNX2, indicating that miR-338-3p activates odontoblast differentiation by targeting RUNX2 [44]. In odontoblast differentiation, Kruppel-like factor 4 (KLF4) represses miR-143, which promotes the expression of miR-145; miR-145 is then able to repress KLF4, consequently forming a feedback loop in the regulation of odontoblast differentiation [45]. In addition, miR-665 targets *Dlx3* mRNA and blocks the expression of *Dlx3* as well as *Runx2*, *Osteorix*, and *Dspp* in mouse odontoblasts [46]. Despite the important role of NF1C in root odontoblast differentiation, the regulatory mechanisms of miRNAs and NF1C in odontoblasts have not been reported to date.

### 3.2. Function of miRNAs in HERS

HERS plays an indispensable role in regulating tooth root development; any malformation of the HERS leads to disturbance in root number, structure, length and abnormalities in periodontal structures [5, 31, 47-49]. By depleting *Alk3* from the dental epithelia, miR-200a and miR-200b were down-regulated, both of which are important for maintaining the epithelial state and prohibiting the epithelial-mesenchymal transition (EMT) by functioning with pro-EMT factors [50]. Consequently, the mutant incisor dental epithelial stem cells differentiated into the root lineage and cementoblast-like cells [51]. Up till now, several cell lines of HERS have been established and several *Cre-LoxP* systems can mediate gene inactivation from HERS cells, however, the functional importance of miRNAs in HERS cells remains largely unknown [2, 31, 52-56].

### 3.3. Function of miRNAs in Periodontal Tissues

Periodontal tissues comprise alveolar bone, periodontal ligaments, gingival, and cementum. The periodontal ligament, alveolar bone, and cementoblasts are derived from dental follicles, a loose connective tissue derived from the ectomesenchyme, which surrounds the developing tooth germ prior to eruption, whereas the gingival tissues are originated from both the oral mucosa and the developing tooth germ [57]. miRNAs play an indispensable role in the homeostasis and pathogenesis of periodontal tissues.

The expression of miR-101 is increased during the osteogenic differentiation of dental follicle cells. When transfected with an miR-101 mimic, SP7 and ALP were increased, indicating that miR-101 sustained the osteogenic differentiation of dental follicle cells [58]. miRNAs have been shown to have functions in the process of periodontitis [59]. The main clinical manifestation of periodontitis is alveolar bone resorption, which involves osteoclasts. miRNAs can regulate the differentiation and function of osteoclasts in periodontal disease [60]. In one study, the authors screened mature miRNA expression in cells treated with tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), a cytokine that promotes alveolar bone resorption, and/or receptor activator for nuclear factor- $\kappa$  B ligand (RANKL), which is expressed in stromal cells, osteoblasts, and T cells and is essential for osteoclast differentiation. During osteoclast formation, the expression of 44 mature miRNAs differed between untreated cells and cells treated with TNF- $\alpha$  plus RANKL; additionally, the expression of 52 mature miRNAs differed upon treatment with RANKL alone, suggesting that

miRNAs may be involved in osteoclast differentiation during the pathogenesis of periodontitis [61]. In a mouse model of periodontitis, miR-146a and several inflammatory cytokines were upregulated compared with that in controls, indicating that miR-146a may directly or indirectly modulate the pathogenesis of periodontitis [62]. The role of miRNAs in osteoclast differentiation and function during alveolar bone loss in periodontitis is critical and has been discussed in several review articles [63, 64].

miRNAs can stimulate the differentiation of periodontal ligament cells (PDLs). In immortalized PDL (I-PDL) lines, nuclear factor kappa-light-chain-enhancer of activated B cells (NF- $\kappa$ B) signaling was believed to regulate bone differentiation. I-PDLs with exogenous miR-146a expression showed attenuation of NF- $\kappa$ B activity and exhibited higher differentiation relative to the controls. Exogenous NF- $\kappa$ B expression decreased the expression of differentiation markers, while the inactivation of endogenous NF- $\kappa$ B increased ALP in I-PDL cells. Additionally, miR-146a was found to promote the differentiation in PDL cells through inhibition of NF- $\kappa$ B signaling [65]. Additionally, in another study, miR-146a was found to play a functional role in regulating a negative feedback loop by decreasing the secretion of pro-inflammatory cytokines and disrupting the Toll-like receptor (TLR) signaling pathway in PDLs stimulated with *Porphyromonas gingivalis* lipopolysaccharide [66]. In mesenchymal stem cells isolated from periodontal ligaments of patients with periodontitis, miR-17 expression was inhibited, and exogenous miR-17 partly reversed the differentiation potential of periodontal ligament tissue-derived mesenchymal stem cells (PDLSCs) in the inflammatory environment. Additionally, the inflammatory microenvironment enhanced the expression of Smad ubiquitin regulatory factor 1 (Smurf1), an important negative regulator of mesenchymal stem cell osteogenic differentiation. Western blotting and 3'-UTR reporter assays identified Smurf1 as a direct target of miR-17 in PDLSCs; thus, inflammatory cytokines caused Smurf1 activation and decreased miR-17 expression, thereby increasing degradation of Smurf1-mediated osteoblast-specific factors [67]. Periodontal ligament-associated protein-1 (PLAP-1/aspurin) regulated the osteogenic differentiation of PDLs, and miR-101 and miR-21 were shown to target PLAP-1, regulating its expression during PDL differentiation [68]. A recent study showed that miR195-5p could regulate osteogenic differentiation of PDLs by directly targeting WNT3A, fibroblast growth factor (FGF) 2 and bone morphogenetic protein receptor (BMPR)-1A, which are important for osteogenic activity and stability under mechanical loading [69].

Previous studies described the occurrence of miRNAs in healthy and inflamed gingival tissues from patients with periodontitis, indicating a role of miRNA in gingival tissues and their pathogenesis [70-72]. With the stimulation of *Porphyromonas gingivalis* (Pg) to human gingival epithelial cells, expressions of several miRNAs (miR-584, miR-203, miR-200, miR-126, miR-155, and miR-210) were altered by targeting different molecules relating to immunity, inflammation and EMT [73-75]. In gingival fibroblasts, deubiquitinase cezanne is a direct target of miR-218 in regulating myofibroblast differentiation in fibroblasts via TGF- $\beta$  [76]. In one study, the author compared the expression of miR-146a and miR-155 in dental pulp (DP), gingival (G) and

**Table 1. miRNAs in tooth root and periodontal tissues.**

Location	MicroRNAs	Target	Origin	Function	Ref.
Dental follicle	miR-98; miR-193 miR-203, miR-27560	(-)	mice	expression	Oommen <i>S., et al.</i> , 2012
	miR101	(-)	human	osteogenic diff. ↑	Klingelhöffer <i>et al.</i> , 2016
Cervical loop	miR-23b, miR-98 miR-148a, miR-193 miR-203, miR-205 miR-218, miR-363 miR-378, miR-27560	(-)	mice	expression	Oommen <i>S., et al.</i> , 2012
DEpSC(P7)	miR200a, miR200b	ZEB1, ZEB2	mice	EMT↓	Yang <i>et al.</i> , 2013
DPaC	miR-34a	NOTCH, TGF-β signaling	human	DPaC diff.↑	Wan, <i>et al.</i> , 2012
	miR-145	KLF4, Osterix	mice	od. diff.↓	Liu, <i>et al.</i> , 2013
DPaSC	miR-34a	NOTCH2, HES1	human	odontogenic, osteogenic diff. ↑	Sun <i>et al.</i> , 2014
DPCs (third molar)	miR-135a	Smad4, Smad5	human	DPC diff. ↓	Song, <i>et al.</i> , 2016
DPSC	miR-720	NANOG	human	stem cell diff. ↑	Hara <i>et al.</i> , 2013
	miR-143, miR-135	MEF2C	human	myogenicdiff.↓	Li <i>et al.</i> , 2015
	miR-424	VEGF, KDR	human	endothelial diff. ↓	Liu <i>et al.</i> , 2014
	miR-152	Sirtuin 7	human	senescence↑, prolif. ↓	Gu <i>et al.</i> , 2016
MDPC-23	miR-27	APC (Wnt/ β-catenin)	mice	od. diff. ↑	Park, <i>et al.</i> , 2014
	miR-663				Kim, <i>et al.</i> , 2014
MDPC6T	miR-338-3p	Runx2	mice	od. diff. ↑	Sun, <i>et al.</i> , 2013
MDPC-23, OD-21, M06-G3	miR-665	Kat6a/Dlx3	mice	od. diff. and maturation↓	Heair <i>et al.</i> , 2015
IPDLC	miR-146a	IRAK1, TRAF6	human	PDL cell diff. ↑	Hung <i>et al.</i> , 2010
P-PDLSC	miR-17	Smurf-1	human	osteogenic diff.: normal condi- tion↓, infla.↑	Liu <i>et al.</i> , 2011
PDLSC	miR-21, miR-101	PLAP-1	human	osteogenic diff. ↓	Li <i>et al.</i> , 2012
PDLSC (under loading)	miR-195-5p	WNT3A, FGF2, BMPRI1A	human	osteogenic diff. ↓	Chang <i>et al.</i> , 2016
PDLSC, DPSC, GSC	miR-218	RUNX2	human	osteogenic diff. ↓	Gay <i>et al.</i> , 2014
PDL progenitor cell	miR-138	osteocalcin	human	cell diff. (under infla.)	Zhou <i>et al.</i> , 2016
PDL fibroblasts	miR-200c	IL-6, IL-8, CCL-5	human	osteogenic diff. ↓	Hong <i>et al.</i> , 2016
PDLSC	miR-21	ACVR2B	human	osteogenic diff. (under stress) ↑	Wei <i>et al.</i> , 2015
gingival epithelial cell (with Pg)	miR-584	LfR	human	anti-infla. ↓	Ouhara <i>et al.</i> , 2014
	miR-203	SOCS3	human	host cell response ↑	Moffatt <i>et al.</i> , 2011
	miR-126, miR-155, miR-210	NFκB, IKKε	human	inflammatory response ↑	Chen <i>et al.</i> , 2016
gingival fibroblast	miR-218	deubiquitinase cezanne	human	myofibroblast diff. ↑	Guo <i>et al.</i> , 2014

\* diff. = differentiation; DPaC: dentalepithelial stem cell; DPSC: dental pulp stem cell; DPC: dental pulp cell; Od.: odontoblast; Prolif.: proliferation; EMT: epithelial mesenchymal transition; MEF2C: myocyte-specific enhancer factor 2C; KLF4: Krüppel-like factor 4; VEGF: vascular endothelial growth factor; KDR: kinase insert domain receptor; APC: adenomatous polyposis coli; IPDLC: immortalized periodontal ligament cell; P-PDLSC: periodontitis-affected periodontal ligament stem cell; infla.: inflammation; LfR: lactoferrin receptor; SOCS3: suppressor of cytokine signaling 3; IKKε: inhibitor-κB kinase ε.

periodontal ligament fibroblasts (PLF) and found that expression of miR-146a and miR-155 are more pronounced in GF than in DPF and PLF, indicating a more prominent role of miR-146a and miR-155 in the regulation GF biological behaviors [77].

### 3.4. Function of miRNAs in Stem Cells from Dental or Periodontal Tissues

Stem cells from dental or periodontal tissues are mainly dental pulp stem cells (DPSC), periodontal ligament stem

cell (PDLSC), stem cells from the apical papilla (SCAP) and gingival stem cell (GSC) [78-81]. In a study conducted in 2014, the author identified that expression of miR-218 was decreased with the differentiation and mineralization of all four stem cell types (bone mesenchymal stem cell, DPSC, GSC and PDLSC) by interacting with Runx2, revealing a regulated role of miRNAs in the differentiation of dental stem cells [82]. Apart from miR-218, several other miRNAs (miR-720, miR-135, miR-143, miR-424 and miR-151) also have different roles in inducing the differentiation of DPSCs, indicating different applications of miRNAs and their inhibitors for therapeutic purposes [83-86]. As for PDLSC, it shares some similarities with osteoblasts and can differentiate into osteoblasts or cementoblasts under certain conditions, it has been selected as potential seeds for tissue regeneration [87]. A previous study found ibandronate, a nitrogen-containing bisphosphonate, could promote the osteogenic differentiation of PDLSCs by regulating the expression of miRNAs (miR125b, miR130a, miR18a, miR19a etc.), which are involved in bone metabolism by regulating the expression of different target genes. Similar investigations on the role of miRNAs in osteogenic differentiation were reported in several papers not only in normal condition but also in inflammation or tissue regeneration [88-92]. Given mechanical force, PDLSCs showed different expression of more than 50 miRNAs (miR21, miR1246 and miR5096 etc.) compared with controls according to the microarray data, and most of the potential target genes are enriched in MAPK and Wnt signaling [93, 94]. Furthermore, in PDLSCs, miRNAs and their inhibitors have potential applications like promoting orthodontic tooth movement, accelerating the healing process of adult stem cell regeneration in smoking people or inducing transdifferentiation into retinal ganglion-like cells for retinal repair purpose [95-97]. In human SCAPs, Notch signaling plays an important role in maintaining of cell differentiation, and miR-34a interacts with Notch signaling to promote both odontogenic and osteogenic differentiation of SCAPs [98].

No studies have yet evaluated the role of miRNAs in the differentiation of cementoblasts. Cementoblasts possess many genetic properties similar to those of osteoblasts and express the same key markers, such as osteopontin, osteonin, and bone sialoprotein [7]. It may be possible to apply the knowledge of osteoblasts to cementoblasts due to their similar biological behaviors. However, further studies are required to confirm this hypothesis.

## CONCLUSION

Tooth root development is a sophisticated process involving interactions between HERS and the adjacent mesenchymal cells, associated with numerous signaling pathways and crosstalk among these pathways. Compared with our knowledge of crown development, little is known about the regulatory mechanisms of tooth root development. To date, the functions of numerous miRNAs have been determined; however, few miRNAs have been shown to participate in root dentin formation, homeostasis, and pathogenesis in periodontal tissues. Moreover, most research explaining miRNA functions have used *in vitro* rather than *in vivo* experiments; accordingly, *in vivo* mouse models may be needed to elucidate the details of the interactions among these cell types.

Further studies are also needed to determine how miRNAs function temporally and interact with other targets, particularly *in vivo*. Besides, deeper understanding of the role of miRNAs in tissue development and pathogenesis will greatly facilitate tooth root regeneration and the use of miRNA target drugs.

## CONSENT FOR PUBLICATION

Not applicable.

## CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

## ACKNOWLEDGEMENTS

This work was supported by Research Grants from the National Nature Science Foundation of China (81030018, 30872900, and 30901680) and the Doctoral Fund of Ministry of Education of China (20120001110043).

## REFERENCES

- [1] Thesleff I. Tooth development. *Dent Update* 1991; 18(9): 382-4, 386-7.
- [2] Zeichner-David M, Oishi K, Su Z, *et al.* Role of Hertwig's epithelial root sheath cells in tooth root development. *Dev Dyn* 2003; 228(4): 651-63.
- [3] Zhang R, Teng Y, Zhu L, *et al.* Odontoblast beta-catenin signaling regulates fenestration of mouse Hertwig's epithelial root sheath. *Sci China Life Sci* 2015; 58(9): 876-81.
- [4] Liu Y, Feng J, Li J, *et al.* An Nfic-hedgehog signaling cascade regulates tooth root development. *Development* 2015; 142(19): 3374-82.
- [5] Yang Y, Ge Y, Chen G, *et al.* Hertwig's epithelial root sheath cells regulate osteogenic differentiation of dental follicle cells through the Wnt pathway. *Bone* 2014; 63: 158-65.
- [6] Torii D, Tsutsui TW, Watanabe N, *et al.* Bone morphogenetic protein 7 induces cementogenic differentiation of human periodontal ligament-derived mesenchymal stem cells. *Odontology* 2016; 104(1): 1-9.
- [7] Zhang R, Yang G, Wu X, Xie J, Yang X, Li T. Disruption of Wnt/beta-catenin signaling in odontoblasts and cementoblasts arrests tooth root development in postnatal mouse teeth. *Int J Biol Sci* 2013. 9(3): 228-36.
- [8] Wang Y, Cox MK, Coricor G, *et al.* Inactivation of Tgfr2 in Osterix-Cre expressing dental mesenchyme disrupts molar root formation. *Dev Biol* 2013; 382(1): 27-37.
- [9] Han XL, Liu M, Voisey A, *et al.* Post-natal effect of overexpressed DKK1 on mandibular molar formation. *J Dent Res* 2011; 90(11): 1312-7.
- [10] Huang X, Xu X, Bringas P Jr, Hung YP, Chai Y. Smad4-Shh-Nfic signaling cascade-mediated epithelial-mesenchymal interaction is crucial in regulating tooth root development. *J Bone Miner Res* 2010. 25(5): 1167-78.
- [11] Cao Z, Liu R, Zhang H, *et al.* Osterix controls cementoblast differentiation through downregulation of Wnt-signaling via enhancing DKK1 expression. *Int J Biol Sci* 2015; 11(3): 335-44.
- [12] Lim WH, Liu B, Cheng D, *et al.* Wnt signaling regulates homeostasis of the periodontal ligament. *J Periodontal Res* 2014; 49(6): 751-9.
- [13] Ambros V. The functions of animal microRNAs. *Nature* 2004; 431(7006): 350-5.
- [14] Kloosterman WP, Plasterk RH. The diverse functions of microRNAs in animal development and disease. *Dev Cell* 2006; 11(4): 441-50.
- [15] Suzuki HI, Yamagata K, Sugimoto K, *et al.* Modulation of microRNA processing by p53. *Nature* 2009; 460(7254): 529-33.
- [16] Cheng CJ, Bahal R, Babar IA, *et al.* MicroRNA silencing for cancer therapy targeted to the tumour microenvironment. *Nature* 2015; 518(7537): 107-10.

- [17] Kasuga H, Fukuyama M, Kitazawa A, *et al.* The microRNA miR-235 couples blast-cell quiescence to the nutritional state. *Nature* 2013; 497(7450): 503-6.
- [18] Powers JT, Tsanov KM, Pearson DS, *et al.* Multiple mechanisms disrupt the let-7 microRNA family in neuroblastoma. *Nature* 2016; 535(7611): 246-51.
- [19] Li X, Carthew RW. A microRNA mediates EGF receptor signaling and promotes photoreceptor differentiation in the *Drosophila* eye. *Cell* 2005; 123(7): 1267-77.
- [20] Hornstein E, Mansfield JH, Yekta S, *et al.* The microRNA miR-196 acts upstream of Hoxb8 and Shh in limb development. *Nature* 2005; 438(7068): 671-4.
- [21] Tu B, Liu S, Yu B, *et al.* miR-203 inhibits the traumatic heterotopic ossification by targeting Runx2. *Cell Death Dis* 2016; 7(10): e2436.
- [22] Jevnaker AM, Osmundsen H. MicroRNA expression profiling of the developing murine molar tooth germ and the developing murine submandibular salivary gland. *Arch Oral Biol* 2008; 53(7): 629-45.
- [23] Thesleff I, Partanen AM, Vainio S. Epithelial-mesenchymal interactions in tooth morphogenesis: the roles of extracellular matrix, growth factors, and cell surface receptors. *J Craniofac Genet Dev Biol* 1991; 11(4): 229-37.
- [24] Huckert M, Stoetzel C, Morkmued S, *et al.* Mutations in the latent TGF-beta binding protein 3 (LTBP3) gene cause brachyolmia with amelogenesis imperfecta. *Hum Mol Genet* 2015; 24(11): 3038-49.
- [25] Guan X, Xu M, Millar SE, *et al.* Beta-catenin is essential for ameloblast movement during enamel development. *Eur J Oral Sci* 2016; 124(3): 221-7.
- [26] Zhu X, Zhao P, Liu Y, *et al.* Intra-epithelial requirement of canonical Wnt signaling for tooth morphogenesis. *J Biol Chem* 2013; 288(17): 12080-9.
- [27] Jheon AH, Prochazkova M, Meng B, *et al.* Inhibition of notch signaling during mouse incisor renewal leads to enamel defects. *J Bone Miner Res* 2016; 31(1): 152-62.
- [28] Huang X, Bringas P Jr, Slavkin HC, Chai Y. Fate of HERS during tooth root development. *Dev Biol* 2009; 334(1): 22-30.
- [29] Huang XF, Chai Y. TGF- $\beta$  signalling and tooth development. *Chin J Dent Res* 2010; 13(1): 7-15.
- [30] Steele-Perkins G, Butz KG, Lyons GE, *et al.* Essential role for NFI-C/CTF transcription-replication factor in tooth root development. *Mol Cell Biol* 2003; 23(3): 1075-84.
- [31] Gao Y, Yang G, Weng T, *et al.* Disruption of Smad4 in odontoblasts causes multiple keratocystic odontogenic tumors and tooth malformation in mice. *Mol Cell Biol* 2009; 29(21): 5941-51.
- [32] Lim WH, Liu B, Hunter DJ, *et al.* Downregulation of Wnt causes root resorption. *Am J Orthod Dentofacial Orthop* 2014; 146(3): 337-45.
- [33] Kim EJ, Lee MJ, Li L, *et al.* Failure of Tooth Formation Mediated by miR-135a Overexpression via BMP Signaling. *J Dent Res* 2014; 93(6): 571-5.
- [34] Oommen S, Otsuka-Tanaka Y, Imam N, *et al.* Distinct roles of microRNAs in epithelium and mesenchyme during tooth development. *Dev Dyn* 2012; 241(9): 1465-72.
- [35] Wan, M, B. Gao, F. Sun, *et al.* microRNA miR-34a regulates cytodifferentiation and targets multi-signaling pathways in human dental papilla cells. *PLoS One* 2012; 7(11): e50090.
- [36] Song Z, Chen LL, Wang RF, *et al.* MicroRNA-135b inhibits odontoblast-like differentiation of human dental pulp cells by regulating Smad5 and Smad4. *Int Endod J*, 2016.
- [37] Park MG, Kim JS, Park SY, *et al.* MicroRNA-27 promotes the differentiation of odontoblastic cell by targeting APC and activating Wnt/beta-catenin signaling. *Gene* 2014; 538(2): 266-72.
- [38] Kim JS, Park MG, Lee SA, *et al.* Downregulation of adenomatous polyposis coli by microRNA-663 promotes odontogenic differentiation through activation of Wnt/beta-catenin signaling. *Biochem Biophys Res Commun* 2014; 446(4): 894-900.
- [39] Bronckers AL, Engelse MA, Cavender A, *et al.* Cell-specific patterns of Cbfa1 mRNA and protein expression in postnatal murine dental tissues. *Mech Dev* 2001; 101(1-2): 255-8.
- [40] Yamashiro T, Aberg T, Levanon D, *et al.* Expression of Runx1, -2 and -3 during tooth, palate and craniofacial bone development. *Mech Dev* 2002; 119 Suppl 1: S107-10.
- [41] Chen S, Rani S, Wu Y, *et al.* Differential regulation of dentin sialophosphoprotein expression by Runx2 during odontoblast cytodifferentiation. *J Biol Chem* 2005; 280(33): 29717-27.
- [42] D'Souza RN, Aberg T, Gaikwad J, *et al.* Cbfa1 is required for epithelial-mesenchymal interactions regulating tooth development in mice. *Development* 1999; 126(13): 2911-20.
- [43] Miyazaki T, Kanatani N, Rokutanda S, *et al.* Inhibition of the terminal differentiation of odontoblasts and their transdifferentiation into osteoblasts in Runx2 transgenic mice. *Arch Histol Cytol* 2008; 71(2): 131-46.
- [44] Sun Q, Liu H, Lin H, *et al.* MicroRNA-338-3p promotes differentiation of mDPC6T into odontoblast-like cells by targeting Runx2. *Mol Cell Biochem* 2013; 377(1-2): 143-9.
- [45] Liu H, Lin H, Zhang L, *et al.* miR-145 and miR-143 regulate odontoblast differentiation through targeting Klf4 and Osx genes in a feedback loop. *J Biol Chem* 2013; 288(13): 9261-71.
- [46] Hear HM, Kemper AG, Roy B, *et al.* MicroRNA 665 regulates dentinogenesis through microRNA-mediated silencing and epigenetic mechanisms. *Mol Cell Biol* 2015; 35(18): 3116-30.
- [47] Luder, H.U. Malformations of the tooth root in humans. *Front Physiol* 2015; 6: 307.
- [48] Ide Y, Nakahara T, Nasu M, *et al.* Cell dynamics in Hertwig's epithelial root sheath and surrounding mesenchyme in mice irradiated to the head. *Oral Dis* 2015; 21(2): 232-9.
- [49] Itaya S, Oka K, Ogata K, *et al.* Hertwig's epithelial root sheath cells contribute to formation of periodontal ligament through epithelial-mesenchymal transition by TGF-beta. *Biomed Res* 2017; 38(1): 61-9.
- [50] Korpala M, Lee ES, Hu G, *et al.* The miR-200 family inhibits epithelial-mesenchymal transition and cancer cell migration by direct targeting of E-cadherin transcriptional repressors ZEB1 and ZEB2. *J Biol Chem* 2008; 283(22): 14910-4.
- [51] Yang Z, Hai B, Qin L, *et al.* Cessation of epithelial Bmp signaling switches the differentiation of crown epithelia to the root lineage in a beta-catenin-dependent manner. *Mol Cell Biol* 2013; 33(23): 4732-44.
- [52] Sonoyama W, Seo BM, Yamaza T, *et al.* Human Hertwig's epithelial root sheath cells play crucial roles in cementum formation. *J Dent Res* 2007; 86(7): 594-9.
- [53] Nam H, Kim J, Park J, *et al.* Expression profile of the stem cell markers in human Hertwig's epithelial root sheath/Epithelial rests of Malassez cells. *Mol Cells* 2011; 31(4): 355-60.
- [54] Arzate, H. Preliminary characterization of epithelial root sheath cells *in vitro*. *Bol Estud Med Biol* 1994; 42(1-4): 27-30.
- [55] Akimoto T, Fujiwara N, Kagiya T, *et al.* Establishment of Hertwig's epithelial root sheath cell line from cells involved in epithelial-mesenchymal transition. *Biochem Biophys Res Commun* 2011; 404(1): 308-12.
- [56] Grachtchouk M, Liu J, Wang A, *et al.* Odontogenic keratocysts arise from quiescent epithelial rests and are associated with deregulated hedgehog signaling in mice and humans. *Am J Pathol* 2006; 169(3): 806-14.
- [57] Fleischmannova J, Matalova E, Sharpe PT, *et al.* Formation of the tooth-bone interface. *J Dent Res* 2010; 89(2): 108-15.
- [58] Klingelhoffer C, Codrin C, Ettl T, *et al.* miRNA-101 supports the osteogenic differentiation in human dental follicle cells. *Arch Oral Biol* 2016; 72: 47-50.
- [59] Gao B, Zheng L. microRNA Expression in Rat Apical Periodontitis Bone Lesion. *Bone Res* 2013; 1(2): 170-85.
- [60] Feng X, Teitelbaum SL. Osteoclasts: New Insights. *Bone Res* 2013; 1(1): 11-26.
- [61] Kagiya, T. and S. Nakamura, Expression profiling of microRNAs in RAW264.7 cells treated with a combination of tumor necrosis factor alpha and RANKL during osteoclast differentiation. *J Periodontol Res* 2013; 48(3): 373-85.
- [62] Nahid MA, Rivera M, Lucas A, *et al.* Polymicrobial infection with periodontal pathogens specifically enhances microRNA miR-146a in ApoE<sup>-/-</sup> mice during experimental periodontal disease. *Infect Immun* 2011; 79(4): 1597-605.
- [63] Kagiya T. MicroRNAs: Potential Biomarkers and Therapeutic Targets for Alveolar Bone Loss in Periodontal Disease. *Int J Mol Sci* 2016; 17(8): pii: E1317.
- [64] Lian J, Du J, Meng S. [A review on microRNAs regulation in osteoclast differentiation and function]. *Hua Xi Kou Qiang Yi Xue Za Zhi* 2015; 33(5): 543-7.
- [65] Hung PS, Chen FC, Kuang SH, *et al.* miR-146a induces differentiation of periodontal ligament cells. *J Dent Res* 2010; 89(3): 252-7.

- [66] Jiang SY, Xue D, Xie YF, *et al.* The negative feedback regulation of microRNA-146a in human periodontal ligament cells after *Porphyromonas gingivalis* lipopolysaccharide stimulation. *Inflamm Res* 2015; 64(6): 441-51.
- [67] Liu Y, Liu W, Hu C, *et al.* MiR-17 modulates osteogenic differentiation through a coherent feed-forward loop in mesenchymal stem cells isolated from periodontal ligaments of patients with periodontitis. *Stem Cells* 2011; 29(11): 1804-16.
- [68] Li C, Yue J, Huang X, *et al.* miR-21 and miR-101 regulate PLAP-1 expression in periodontal ligament cells. *Mol Med Rep* 2012; 5(5): 1340-6.
- [69] Chang M, Lin H, Fu H, *et al.* MicroRNA-195-5p regulates osteogenic differentiation of periodontal ligament cells under mechanical loading. *J Cell Physiol* 2017. [Epub ahead of print].
- [70] Xie YF, Shu R, Jiang SY, *et al.* Comparison of microRNA profiles of human periodontal diseased and healthy gingival tissues. *Int J Oral Sci* 2011; 3(3): 125-34.
- [71] Stoecklin-Wasmer C, Guarnieri P, Celenti R, *et al.* MicroRNAs and their target genes in gingival tissues. *J Dent Res* 2012; 91(10): 934-40.
- [72] Ogata Y, Matsui S, Kato A, *et al.* MicroRNA expression in inflamed and noninflamed gingival tissues from Japanese patients. *J Oral Sci* 2014; 56(4): 253-60.
- [73] Ouhara K, Savitri IJ, Fujita T, *et al.* miR-584 expressed in human gingival epithelial cells is induced by *Porphyromonas gingivalis* stimulation and regulates interleukin-8 production via lactoferrin receptor. *J Periodontol* 2014; 85(6): e198-204.
- [74] Moffatt CE, Lamont RJ. *Porphyromonas gingivalis* induction of microRNA-203 expression controls suppressor of cytokine signaling 3 in gingival epithelial cells. *Infect Immun* 2011; 79(7): 2632-7.
- [75] Chen SC, Constantinides C, Kebschull M, *et al.* MicroRNAs Regulate Cytokine Responses in Gingival Epithelial Cells. *Infect Immun* 2016; 84(12): 3282-9.
- [76] Guo F, Carter DE, Leask A. miR-218 regulates focal adhesion kinase-dependent TGFbeta signaling in fibroblasts. *Mol Biol Cell* 2014; 25(7): 1151-8.
- [77] Sipert CR, Morandini AC, Dionisio TJ, *et al.* MicroRNA-146a and microRNA-155 show tissue-dependent expression in dental pulp, gingival and periodontal ligament fibroblasts *in vitro*. *J Oral Sci* 2014; 56(2): 157-64.
- [78] Gronthos S, Mankani M, Brahimi J, *et al.* Postnatal human dental pulp stem cells (DPSCs) *in vitro* and *in vivo*. *Proc Natl Acad Sci USA* 2000; 97(25): 13625-30.
- [79] Ikeda, E. and T. Tsuji, Growing bioengineered teeth from single cells: potential for dental regenerative medicine. *Expert Opin Biol Ther* 2008; 8(6): 735-44.
- [80] Gould TR, Melcher AH, Brunette DM. Location of progenitor cells in periodontal ligament of mouse molar stimulated by wounding. *Anat Rec* 1977; 188(2): 133-41.
- [81] Gronthos S, Mrozik K, Shi S, *et al.* Ovine periodontal ligament stem cells: isolation, characterization, and differentiation potential. *Calcif Tissue Int* 2006; 79(5): 310-7.
- [82] Gay I, Cavender A, Peto D, *et al.* Differentiation of human dental stem cells reveals a role for microRNA-218. *J Periodontol Res* 2014; 49(1): 110-20.
- [83] Hara ES, Ono M, Eguchi T, *et al.* miRNA-720 controls stem cell phenotype, proliferation and differentiation of human dental pulp cells. *PLoS One* 2013; 8(12): e83545.
- [84] Li D, Deng T, Li H, *et al.* MiR-143 and miR-135 inhibitors treatment induces skeletal myogenic differentiation of human adult dental pulp stem cells. *Arch Oral Biol* 2015; 60(11): 1613-7.
- [85] Liu W, Gong Q, Ling J, *et al.* Role of miR-424 on angiogenic potential in human dental pulp cells. *J Endod* 2014; 40(1): 76-82.
- [86] Gu S, Ran S, Liu B, *et al.* miR-152 induces human dental pulp stem cell senescence by inhibiting SIRT7 expression. *FEBS Lett* 2016; 590(8): 1123-31.
- [87] Seo BM, Miura M, Gronthos S, *et al.* Investigation of multipotent postnatal stem cells from human periodontal ligament. *Lancet* 2004; 364(9429): 149-55.
- [88] Chao D, Yan W, Kun Y, *et al.* [Effect of microRNA-17 on osteogenic differentiation of advanced glycation end products-stimulated human periodontal ligament stem cells]. *Hua Xi Kou Qiang Yi Xue Za Zhi* 2015; 33(1): 21-4.
- [89] Zhou X, Luan X, Chen Z, *et al.* MicroRNA-138 Inhibits Periodontal Progenitor Differentiation under Inflammatory Conditions. *J Dent Res* 2016; 95(2): 230-7.
- [90] Hong L, Sharp T, Khorsand B, *et al.* MicroRNA-200c represses IL-6, IL-8, and CCL-5 expression and enhances osteogenic differentiation. *PLoS One* 2016; 11(8): e0160915.
- [91] Diomedea F, Merciaro I, Martinotti S, *et al.* miR-2861 is involved in osteogenic commitment of human periodontal ligament stem cells grown onto 3D scaffold. *J Biol Regul Homeost Agents* 2016; 30(4): 1009-18.
- [92] Hao Y, Ge Y, Li J, *et al.* Identification of MicroRNAs by microarray analysis and prediction of target genes involved in the osteogenic differentiation of human periodontal ligament stem cells. *J Periodontol* 2017; 1-11.
- [93] Wei F, Liu D, Feng C, *et al.* microRNA-21 mediates stretch-induced osteogenic differentiation in human periodontal ligament stem cells. *Stem Cells Dev* 2015; 24(3): 312-9.
- [94] Wei FL, Wang JH, Ding G, *et al.* Mechanical force-induced specific MicroRNA expression in human periodontal ligament stem cells. *Cells Tissues Organs* 2014; 199(5-6): 353-63.
- [95] Chen N, Sui BD, Hu CH, *et al.* microRNA-21 Contributes to Orthodontic Tooth Movement. *J Dent Res* 2016; 95(12): 1425-33.
- [96] Ng TK, Carballosa CM, Pelaez D, *et al.* Nicotine alters MicroRNA expression and hinders human adult stem cell regenerative potential. *Stem Cells Dev* 2013; 22(5): 781-90.
- [97] Ng TK, Yung JS, Choy KW, *et al.* Transdifferentiation of periodontal ligament-derived stem cells into retinal ganglion-like cells and its microRNA signature. *Sci Rep* 2015; 5: 16429.
- [98] Sun F, Wan M, Xu X, *et al.* Crosstalk between miR-34a and Notch Signaling Promotes Differentiation in Apical Papilla Stem Cells (SCAPs). *J Dent Res* 2014; 93(6): 589-95.