

# Scientific Article

## Risk Factors for Severe Early Childhood Caries in Children Younger Than 4 Years Old in Beijing, China

Man Qin, DDS, PhD<sup>1</sup> • Jing Li, DDS, MSc<sup>2</sup> • Sun Zhang, DDS, MSc<sup>3</sup> • Wenli Ma, DDS, MSc<sup>4</sup>

**Abstract:** *Purpose:* The purpose of this study was to compare cariogenic factors and acidogenic ability of bacteria between severe early childhood caries (S-ECC) and caries-free children in Beijing, China. *Methods:* The study consisted of 117 S-ECC children and 129 caries-free children < 4 years old. A questionnaire was designed to collect background information, feeding habits, and oral hygiene practices. Dental plaque samples were collected to test acidogenic ability of bacteria. *Results:* Compared with the mothers of caries-free children, those of S-ECC children had a lower education level and poorer knowledge of oral hygiene ( $P < .05$ ). Night-feeding and eating sweets several times a day were significantly more common in S-ECC children than in caries-free children ( $P < .001$ ). Forty-six S-ECC children but only 2 caries-free children received prechewed food ( $P < .001$ ). The results of the Cariostat test showed that the majority of the caries-free children (81%) were at a low risk level, and most of the S-ECC children (78%) were at a high risk level ( $P < .001$ ). *Conclusions:* There was strong relationship between high acidogenic ability of bacteria and S-ECC. A lower maternal education level, poor knowledge of oral hygiene, night-feeding, and excessive sugar intake were important contributors to the development of S-ECC. (*Pediatr Dent* 2008;30:122-8) Received March 20, 2007 / Last Revision July 29, 2007 / Revision Accepted August 13, 2007.

KEYWORDS: SEVERE EARLY CHILDHOOD CARIES (S-ECC), ORAL HEALTH BEHAVIORS, ACIDOGENIC ABILITY OF BACTERIA

The concepts of early childhood caries (ECC) and severe early childhood caries (S-ECC) have been used for nearly 10 years to describe caries status present in children younger than 6 years old.<sup>1,2</sup> S-ECC has been defined in children younger than 3 years old as any sign of smooth surface caries. It is defined for children 3 to 5 years as 1 or more cavitated, missing (due to caries), or filled surfaces in primary maxillary teeth or as a decayed, missing, or filled surface with a score  $\geq 4$  at age 3,  $\geq 5$  at age 4, or  $\geq 6$  at age 5.<sup>2</sup>

ECC is a chronic, transmissible, infectious disease with a complex and multifactorial etiology. Factors attributed to the etiology of ECC include: excessive bottle-feeding with sugar-containing liquids; breast-feeding on demand and/or falling asleep while feeding; and nursing beyond the recommended age for weaning.<sup>3,4</sup> Other factors associated with ECC include: genetic predisposition; parental education; and nutritional, environmental, socioeconomic, and parental style factors.<sup>5,6</sup> Although the risk factors for ECC have been well reported in

many countries, the status and health behaviors associated with ECC in Chinese children, particularly those younger than 5 years old, are not yet clear. One reason is that children < 5 years old are not included in the National Oral Health Survey.

Furthermore, an early cariogenic bacterial infection is one of the important predisposing factors of ECC.<sup>7</sup> As mutans streptococci (MS) metabolize nutrients, they produce acidic byproducts that penetrate the enamel pellicle. Lactobacilli present in dental plaque can also colonize damaged areas, producing more acid byproducts and further damaging the interior of the tooth.<sup>8</sup> It is reported that MS and lactobacilli are closely related to incipient caries and to caries progression.<sup>9-10</sup> A caries activity test is designed to test the acidogenic ability of dental plaque, including MS and lactobacilli.

The Cariostat test medium (Cariostat, Sankin Co, Tokyo, Japan) is a commercial product containing highly concentrated sucrose and 2 kinds of pH indicators to show the continuous pH decrease of the test medium caused by bacteria, including MS and lactobacilli, in plaque samples.<sup>11</sup> Cariostat scores of 0, 1, 2, and 3 correspond to the pH values of 5.8-7.2,  $5.4 \pm 0.3$ ,  $4.7 \pm 0.3$ , and  $< 4.4$ , respectively. The test does not measure bacteria directly, but it does measure their acidogenic ability, which indicates the infection levels of MS and lactobacilli in the plaque.<sup>12</sup> The test has high sensitivity and specificity and is recognized as a screening method for primary caries.<sup>13</sup>

<sup>1</sup>Dr. Qin is professor and vice-director, <sup>2</sup>Dr. Li is assistant clinical professor, <sup>3</sup>Dr. Zhang is associate professor, and <sup>4</sup>Dr. Ma is associate clinical professor, all in the Department of Pediatric Dentistry, Peking University School and Hospital of Stomatology, Beijing, China.

Correspond with Dr. Qin at qinman@gmail.com.

The aims of the present study were to:

1. characterize and compare behavioral and environmental factors potentially associated with caries in children with S-ECC and caries-free children <4 years old in Beijing, China; and
2. compare the acidogenic ability of dental plaque collected from S-ECC children and caries-free children by using the Cariostat test.

## Methods

**Subjects.** Five hundred fourteen children <4 years old, recruited from the Pediatric Dental Clinic of Peking University Stomatological Hospital, Beijing, China, and 5 kindergarteners located within 5 km of the hospital were invited to participate in the study. A total of 117 children with >5 decayed teeth were placed into the S-ECC group, and 129 caries-free children were placed in the control group. Children with discernible enamel hypoplasia were not included in the study. The ethical approval of this study was obtained from the Human Research Ethics Committee of Peking University Medical Science Centre, Beijing, China.

**Parental interview.** The study was explained to the parents and legal guardians, and an informed consent was obtained from those who agreed to answer a detailed questionnaire and allow their children to undergo a dental examination and dental plaque collection. No more than 20 parents were interviewed at one time. The content of the questionnaire was described to the parents, and only the mothers were asked to complete the questionnaire in the interview room. The questionnaire was designed to collect information regarding the mother's education and knowledge of basic dental care and her child's birth weight, feeding practices, dental care, and hygiene behaviors.

**Dental examination.** A dental examination was performed by 2 pediatric dentists using a mirror and probe at the dental chair in the clinic with a knee-to-knee approach for the kindergarteners. The examiners were calibrated for caries detection by examining 20 children younger than <4 years old. The degree of agreement was assessed with the kappa statistic. Both examiners decided if a child was excluded due to enamel hypoplasia. The teeth were cleaned, and no radiographs were taken. Decayed, missing, and filled primary teeth (dmft) and surfaces (dmfs) and decayed teeth (dt) and surfaces (ds) were assessed according to the dental caries diagnostic criteria of the World Health Organization (WHO).<sup>14</sup> Restored teeth with recurrent caries were considered decayed.

**Cariostat test.** The plaque samples were collected from primary maxillary and mandibular buccal surfaces using a sterile cotton swab before cleaning the teeth. The swab was scrubbed across the tooth surface 3 or 4 times with a wiping movement. The cotton swab was placed into an ampule containing 2 mL of the Cariostat

test medium. The sample was incubated at 37°C for 48 hours. The Cariostat score was evaluated by referencing the sample with 4 standard color tubes (provided by the manufacturer). In this study, a modified scale was used, where the interval between 0 and 1.0, 1.0 and 2.0, and 2.0 and 3.0 were divided into halves.

**Statistical analysis.** The data were analyzed by the Working Group of Medical Statistics, Peking University Medical College, Beijing, China. Data were divided into S-ECC and caries-free groups and calculated by descriptive statistics using SPSS v. 10.0 software for Windows (SPSS, Inc, Chicago, Ill). Pearson's chi-square test was used to analyze the difference in psychosocial, behavioral, and environmental factors between S-ECC children and caries-free children. Fisher's exact test was used when the *t* value was within the range of 1% to 5%. The Cariostat test scores of the S-ECC children and caries-free children were compared using the Mann-Whitney test. Differences were regarded as statistically significant if  $P < .05$ .

## Results

**Clinical findings.** The kappa score for the interexaminer calibration was 0.91, indicating very good examiner agreement. The caries experience of the S-ECC children is summarized in Table 1. The mean dmft and dmfs were similar to the mean dt and ds, respectively.

Table 1. CARIES SCORES OF CHINESE CHILDREN IN THE SEVERE EARLY-CHILDHOOD CARIES GROUP

| Age (mos)  | N   | dmft±SD | dt±SD   | dmfs±SD  | ds±SD    |
|------------|-----|---------|---------|----------|----------|
| <24        | 23  | 9.2±3.6 | 8.8±3.6 | 15.5±7.7 | 14.8±7.7 |
| 24-35      | 67  | 9.3±3.6 | 8.8±3.6 | 15.6±7.8 | 15.0±7.8 |
| 36-48      | 27  | 9.2±3.6 | 8.8±3.6 | 15.5±7.8 | 14.9±7.7 |
| Total      |     |         |         |          |          |
| 29.84±7.69 | 117 | 9.2±3.6 | 8.8±3.6 | 15.5±7.7 | 14.9±7.7 |

**Basic information of children and parents.** The general information of the children and their parents is presented in Table 2. The differences in birth weight between S-ECC children and caries-free children were not significant ( $P = .156$ ). The mother's education level for caries-free children was higher than that for S-ECC children ( $P = .023$ ). More S-ECC children were cared for by grandparents or a nursery maid than caries-free children ( $P = .041$ ). Thirty-nine percent of S-ECC children had received prechewed food, of which one third received prechewed food from their grandparents and two thirds had received prechewed food from their parents. Only 2 caries-free children, however, had received prechewed food from their parents. This difference was statistically significant ( $P < .001$ ). Among the 46 children who received prechewed food, 43 had

**Table 2. INFORMATION ABOUT CHINESE CHILDREN AND THEIR PARENTS ENROLLED IN THE STUDY**

|                                    | Severe early childhood caries group* | Caries-free group† | P-value of chi-square test |
|------------------------------------|--------------------------------------|--------------------|----------------------------|
|                                    | N (%)                                | N (%)              |                            |
| Female                             | 53 (45)                              | 60 (47)            |                            |
| Birth weight                       |                                      |                    | .156                       |
| <2,500 g                           | 5 (4)                                | 3 (2)              |                            |
| 2,500-3,000 g                      | 11 (9)                               | 24 (19)            |                            |
| >3,000 g                           | 101 (86)                             | 102 (79)           |                            |
| Mother's education level‡          |                                      |                    | .023                       |
| Elementary                         | 7 (6)                                | 2 (2)              |                            |
| High school                        | 34 (29)                              | 21 (16)            |                            |
| College and higher                 | 76 (65)                              | 106 (82)           |                            |
| Person(s) taking care of the child |                                      |                    | .041                       |
| Parent(s)                          | 39 (33)                              | 49 (38)            |                            |
| Parent(s) and grandparent(s)       | 34 (29)                              | 53 (41)            |                            |
| Grandparent(s)                     | 24 (21)                              | 15 (11)            |                            |
| Nursery maid(s)                    | 20 (17)                              | 12 (10)            |                            |
| Fed with prechewed food            |                                      |                    | <.001                      |
| No                                 | 71 (61)                              | 127 (99)           |                            |
| Yes                                | 46 (39)                              | 2 (2)              |                            |
| Fed by parent(s)                   | 35 (30)                              | 2 (2)              |                            |
| Fed by grandparent(s)              | 11 (9)                               | 0                  |                            |
| Fed by others                      | 0                                    | 0                  |                            |
| Reason for first dental visit      |                                      |                    | <.001                      |
| Routine dental examination         | 2 (2)                                | 32 (25)            |                            |
| Dental caries                      | 86 (74)                              | 0                  |                            |
| Toothache or/and gingival abscess  | 25 (21)                              | 0                  |                            |
| Dental trauma                      | 4 (3)                                | 3 (2)              |                            |
| Never see a dentist                | 0                                    | 94 (73)            |                            |

\* Average age in mos±(SD)=26.6±8.9.

† Average age in mos±(SD)=41.6±4.5.

‡ Fisher's exact test was used because the T value was within the 1%-5% range in Pearson's chi-square.

mothers with a high school or lower education level. Ninety-four of 129 caries-free children never saw a dentist before this dental examination, while 32 of them received routine dental examinations. Most S-ECC children (95%) did not visit a dentist until either serious caries decay was observed or they suffered a toothache, while only 2 children had their first dental visit as a routine dental examination ( $P<.001$ ).

**Table 3. COMPARISON OF FEEDING HABITS BETWEEN CHINESE CHILDREN WITH SEVERE EARLY CHILDHOOD CARIES (S-ECC) AND CARIES-FREE CHILDREN**

|  | S-ECC group N (%) | Caries-free group N (%) | P-value of chi-square test |
|--|-------------------|-------------------------|----------------------------|
| Feeding habit before 6 mos old                             |                   |                         | .002                       |
| Breast-feeding   | 65 (56)           | 65 (50)                 |                            |
| Milk without sugar   | 4 (3)             | 23 (18)                 |                            |
| Milk with sugar  | 48 (41)           | 41 (32)                 |                            |
| Feeding habit for 6- to 12-mos-olds                        |                   |                         |                            |
| Breast-feeding   | 39 (33)           | 25 (19)                 |                            |
| Milk without sugar   | 8 (7)             | 39 (30)                 |                            |
| Milk with sugar  | 70 (60)           | 65 (50)                 |                            |
| Sleeping while feeding after 12 mos old                    |                   |                         | <.001                      |
| No   | 40 (34)           | 112 (87)                |                            |
| Yes  | 77 (66)           | 17 (13)                 |                            |
| Milk without sugar   | 65 (55)           | 4 (3)                   |                            |
| Milk with sugar  | 12 (11)           | 13 (10)                 |                            |
| Drinking habit   |                   |                         | <.001                      |
| Water only   | 12 (10)           | 19 (14)                 |                            |
| Mainly water, a small amount of sugar-containing beverages | 53 (45)           | 79 (61)                 |                            |
| Mainly sugar-containing beverages, a little water          | 15 (13)           | 17 (14)                 |                            |
| Sugar-containing beverages only                            | 37 (32)           | 14 (11)                 |                            |
| Bottle-feeding by age (mos)                                |                   |                         | .23                        |
| No bottle-feeding  | 14 (12)           | 10 (8)                  |                            |
| <12  | 0                 | 2 (2)                   |                            |
| 12-18  | 7 (6)             | 18 (14)                 |                            |
| 18-24  | 19 (16)           | 11 (8)                  |                            |
| >24  | 77 (66)           | 88 (68)                 |                            |
| Frequency of eating sweets or snacks every day             |                   |                         | <.001                      |
| Never  | 6 (5)             | 4 (3)                   |                            |
| Rarely   | 17 (15)           | 53 (41)                 |                            |
| 1-2x/day   | 25 (21)           | 52 (40)                 |                            |
| >2x/day  | 69 (59)           | 20 (16)                 |                            |

**Feeding habits and dental health behaviors.** Information on feeding habits and dental health behaviors of the children are presented in Tables 3 and 4. The feeding habits from birth to 12 months of age were significantly different between S-ECC children and caries-free children ( $P<.001$ ). The S-ECC children were fed with milk containing more sugar than were the caries-free children, though S-ECC children seemed to receive more breast-feeding. Compared with 66% of S-ECC children who drank milk at night, 87% of caries-free children did not drink milk at night ( $P<.001$ ).

Table 4. COMPARISON OF DENTAL HEALTH BEHAVIORS BETWEEN CHINESE CHILDREN WITH SEVERE EARLY CHILDHOOD CARIES (S-ECC) AND CARIES-FREE CHILDREN

|   | S-ECC group<br>N (%) | Caries-free group<br>N (%) | P-value of<br>chi-square<br>test |
|---|----------------------|----------------------------|----------------------------------|
| Mother's knowledge about when to begin cleaning her child's mouth or brushing the child's teeth |                      |                            |                                  |
|   |                      |                            | <.001                            |
| Before the first tooth erupted  | 8 (7)                | 6 (5)                      |                                  |
| By the time the first tooth erupted   | 21 (18)              | 25 (19)                    |                                  |
| After all primary teeth erupted   | 49 (41)              | 84 (65)                    |                                  |
| Do not know/no idea   | 39 (34)              | 14 (11)                    |                                  |
| Frequency of toothbrushing  |                      |                            |                                  |
|   |                      |                            | 6.58                             |
| ≥1x/day   | 43 (37)              | 38 (29)                    |                                  |
| ≥3x/week  | 14 (12)              | 28 (22)                    |                                  |
| <3x/week  | 33 (28)              | 44 (34)                    |                                  |
| Never   | 27 (23)              | 19 (15)                    |                                  |
| How to clean child's mouth  |                      |                            |                                  |
|   |                      |                            | .87                              |
| Parent(s) brush for the child   | 54 (46)              | 40 (31)                    |                                  |
| Brushing by the child   | 10 (9)               | 44 (35)                    |                                  |
| Rinse with water only   | 27 (23)              | 29 (22)                    |                                  |
| None  | 26 (22)              | 16 (12)                    |                                  |

Table 5. COMPARISON OF THE CARIOSTAT SCORE BETWEEN CHINESE CHILDREN WITH SEVERE EARLY CHILDHOOD CARIES (S-ECC) AND CARIES-FREE CHILDREN

| Cariostat score | S-ECC GROUP |     | Caries-free group |     | Mann-Whitney test |         |
|-----------------|-------------|-----|-------------------|-----|-------------------|---------|
|                 | N           | %   | N                 | %   | Z                 | P-value |
| 0-0.5           | 0           | 0   | 43                | 33  |                   |         |
| 1.0             | 6           | 5   | 62                | 48  |                   |         |
| 1.5             | 20          | 17  | 24                | 19  |                   |         |
| 2.0             | 40          | 34  | 0                 | 0   |                   |         |
| 2.5-3.0         | 51          | 44  | 0                 | 0   |                   |         |
| Total           | 117         | 100 | 129               | 100 | 12.66             | <.001   |

Sixty-six percent of S-ECC children and 68% of caries-free children continued bottle-feeding beyond 24 months old. The difference in bottle-feeding age between S-ECC children and caries-free children was not significant ( $P=.231$ ). Seventy-five percent of caries-free children drank only water daily, or primarily water and a small amount of sugar-containing beverages, while 44% of S-ECC children drank only sugar-containing beverages or sugar-containing beverages and a little water ( $P<.001$ ). Eighty-four percent of caries-free children had sweets less than twice a day, while 59% of S-ECC children had sweets several times a day ( $P<.001$ ). In addition, the results showed that the mothers of S-ECC children had a poor knowledge of their children's oral hygiene. Mothers of S-ECC children also did not know when to begin cleaning their children's teeth, compared to 89% of mothers of caries-free children who knew that they should begin cleaning

their children's mouths or brushing their children's teeth no later than when all primary teeth erupt. The difference was statistically significant ( $P<.001$ ). No significant difference was observed in the frequency of tooth-brushing and the method of mouth cleaning between S-ECC children and caries-free children.

**Acidogenic ability of dental bacteria.** The results of the Cariostat test are shown in Table 5. The difference between S-ECC children and caries-free children was statistically significant ( $P<.001$ ). Most caries-free children (81%) were at a low risk level (value  $\leq 1.0$ ,  $\text{pH} > 5.5$ ), and none were at a high risk level (value  $\geq 2.0$ ,  $\text{pH} < 5.0$ ). Alternatively, most of the S-ECC children (78%) were at a high risk level. Seventeen percent of S-ECC children and 19% of caries-free children were at a medium risk level (value=1.5,  $\text{pH}=5.5-5.0$ ).

## Discussion

In this study, dt was the major component of the dmft score of the S-ECC children. Although these S-ECC children had a very serious dental condition, most (95%) did not visit a dentist until serious decay was observed or they suffered a toothache. Their parents may have been unaware of the need for treatment or perhaps felt that their children were too young to attend a dental clinic until pain was experienced.<sup>15</sup> The same observations were reported by Wong et al<sup>16,17</sup> among 5- to 6-year-old children in China in 1997 and 2001.

Poor knowledge of oral hygiene and dental care by the parents was also related to the development of S-ECC.<sup>18-20</sup> One third of the mothers of S-ECC children did not know when they should begin to take care of their children's teeth, while most (89%) of the caries-free children's mothers knew that they should clean their children's mouths or brush their teeth no later than the eruption of all primary teeth. Although the mothers of the caries-free children had a better understanding of oral hygiene, no relationship was found between oral hygiene habits and dental caries in this study.

Because of the difficulties in brushing the children's teeth, most mothers let their children brush their teeth themselves, with less than half of both the S-ECC children and caries-free children receiving effective dental care (toothbrushing by caregivers everyday). Riedy et al<sup>21</sup> reported a similar phenomenon and argued that expectations about tooth cleaning behavior should be realistic, stressing persistence and the encouragement of children to cooperate with repeated, regular practice. In addition, for parents with very young children, it might be helpful to show how tooth brushing can be a kind of game. The fact that more S-ECC children were cared for by grandparents or a nursery maid than were caries-free children ( $P=.041$ ) reflected a poor knowledge of oral hygiene among grandparents and their low opinion of primary teeth. In China,

some adults believe that a child under school age is too young for toothbrushing. They believe that primary teeth are transitional and that the decay of primary teeth will not affect permanent teeth. Therefore, they believe that it is unnecessary to care for primary teeth. It is imperative to improve public knowledge of oral hygiene to correct traditional norms and to promote dental care for young children in the fight against ECC in China.

It is well recognized that a low education level among parents is a risk factor associated with ECC.<sup>2,6</sup> The situation of ECC in mainland China is unclear, since few studies have been undertaken to evaluate the epidemiology and etiology of ECC. Bu et al<sup>22</sup> investigated 551 preschool children in Guangzhou, a city in south China. Their results showed that a parent's low education level and young age were risk factors for children to suffer caries. Huang,<sup>23</sup> however, investigated 890 preschool children in Guangxi, a province in southwest China, and found an opposite result. His study showed that the mother's education level did not significantly associate with ECC. In his study, 80% of children in high-income families suffered ECC, compared to 48% of children in low-income families. In this study, 65% of mothers of S-ECC children and 82% of mothers of caries-free children claimed to have a college or higher education. It seemed that mothers of both S-ECC children and caries-free children belonged to a population of higher education because less than 10% of people receive college or higher education in China. Nevertheless, the education level of S-ECC children's mothers was significantly lower than that for caries-free children, and a mother's lower education level is still a risk factor associated with their children suffering caries.

Improper feeding habits were thought to be important attributive factors for S-ECC. Frequent consumption of liquids containing fermentable carbohydrates (juice, milk, infant formula, soda) was reported to increase the risk of caries due to prolonged contact between sugars in the consumed liquid and cariogenic bacteria on the susceptible teeth.<sup>24</sup> Parents are encouraged to use bottle-feeding appropriately, have infants drink from a cup as they approach their first birthday, and wean children from the bottle at 12 to 14 months of age.<sup>25</sup> This recommendation is based on the belief that prolonged use of the baby bottle is an underlying cause of caries among young children. Nevertheless, some studies indicate that the relationship between prolonged use of the baby bottle and caries is weak, and that feeding at night and misuse of sugar seem to be the major contributors to the development of S-ECC.<sup>26-29</sup> Similar results were observed in this study.

Giving a child prechewed food is thought to increase the risk for dental caries because it could transmit cariogenic microorganisms directly to a child. This feeding behavior was rarely mentioned in previous studies. Feeding prechewed food is sometimes seen in China, particularly in the traditional practices of elderly Chinese. In this study, 39% of S-ECC children received prechewed food, compared to 2% of caries-free children. Forty-three of 46 S-ECC children who received prechewed food belonged to mothers with a lower education level. The acidogenic ability of dental plaque of the 46 S-ECC children

was not significantly different from the other S-ECC children who did not receive prechewed food. The relationship between receiving prechewed food, the level of micro-organisms, and the acidogenic ability of dental plaque in S-ECC children was not clear because the prechewed food behavior is random; it is very difficult to determine its frequency and duration.

Low birth weight is reported to be a risk factor related to S-ECC, since low birth weight could result in a higher prevalence of enamel hypoplasia.<sup>30,31</sup> In this study, birth weight was not significantly different between S-ECC children and caries-free children. This might be a result of the fact that children suffering discernible enamel hypoplasia were excluded from the study. Fluoride exposure is an important factor related to ECC. Fluoride has 3 principle topical mechanisms of action: (1) inhibiting bacterial metabolism; (2) inhibiting demineralization; and (3) enhancing remineralization.<sup>32,33</sup> In this study, the fluoride exposure of the 2 groups was at the same baseline. Fluoride concentration in the drinking water in Beijing is below 0.4 ppm. Also, fluoride supplements are not available.

The topical application of fluoride is not carried out until "the first dental examination" provided by the Health Office for Women and Infants, a local governmental health unit. In fact, the dental examination in the study was part of their "first dental examination." In addition, only 6 of 117 S-ECC children had received dental treatment, but no topical application of fluoride was found in their dental records. Fluoridated toothpaste is available in Beijing stores and is recommended for children older than 3 to 4 years old. In this study, no mother declared giving her child fluoridated toothpaste regularly due to the risk of swallowing fluoridated toothpaste.

At least 3 categories of risk factors are associated with caries development: (1) microorganisms; (2) substrate/oral environment; and (3) host/teeth. Several lines of evidence strongly suggest that the first step in the development of ECC is MS infection. MS initiate the colonization and demineralization of tooth enamel; dietary factors contribute to the severity of ECC, but do not initiate the process.<sup>32-36</sup> Ansai et al<sup>37</sup> studied the relationship between dental caries experience (dfs) of 4- and 5-year-old Japanese children and the results of caries activity tests. These included salivary levels of MS (the Mucount test) and the acidogenic ability of dental plaque bacteria (the Cariostat test). They reported a significant positive correlation between the results of the Mucount test and the Cariostat tests in the total group of subjects.

In this study, a strong association between microorganisms and S-ECC was shown using the Cariostat test. Most caries-free children (81%) were at a low risk level (pH>5.5), and none was at a high-risk level (pH<5.0). Most of the S-ECC children (78%), however, were at a high risk level. In both groups, nearly one fifth of the children showed a medium level of acidogenic ability of dental bacteria. Six children with more than 8 decayed teeth were observed at the critical risk level of acidogenic ability of dental bacteria. The common variables among these 6 subjects were being male, prolonged bottle-feeding beyond 24 months, and sweet drinks at night.



It might be that microorganisms were not the sole cariogenic factor at the time, but that the host/teeth and dietary factors contributed together to the etiology of caries. Further studies should observe the caries risk in caries-free children with a medium level of acidogenic ability of dental bacteria.

### Conclusions

Based on this study's results, the following conclusions can be made:

1. There was strong relationship between high acidogenic ability of bacteria and severe early childhood caries (S-ECC).
2. A lower maternal education level, poor knowledge of oral hygiene, night-feeding, and excessive sugar intake were important contributors to the development of S-ECC in young children. Giving child prechewed food should be considered a potential caries risk factor among children in Beijing, China.
3. Health promotion strategies that target new and expectant mothers, and that promote a more positive attitude towards oral health in China should be established.

### Acknowledgments

The authors would like to thank Professor Tsutomu Shimono, Department of Pediatric Dentistry, Okayama University, Okayama City, Japan) for his help in the Cariostat test and Ms. Xueying Li for her assistance with the statistics. The Fund of Beijing Medical Research and Development, Grant No.2002-3035, supported this study.

### References

1. Drury TF, Horowitz AM, Ismail AI, Maertens MP, Rozier RG, Selwitz RH. Diagnosing and reporting early children caries for research purposes. A report of a workshop sponsored by the National Institute of Dental and Craniofacial Research, the Health Resources and Services Administration, and the Health Care Financing Administration. *J Public Health Dent* 1999;59:192-7.
2. Veerkamp JSJ, Weerheijm KL. Nursing-bottle caries. The importance of a developmental perspective. *J Dent Child* 1995;62:381-6.
3. Hallett KB, O'Rourke PK. Early childhood caries and infant feeding practice. *Community Dent Health* 2002; 19:237-42.
4. Hallonsten AL, Wendt LK, Mejare I, et al. Dental caries and prolonged breast-feeding in 18-month-old Swedish children. *Int J Paediatr Dent* 1995;5:149-55.
5. Febres C, Echeverri EA, Keene HJ. Parental awareness, habits, and social factors and relationship to baby bottle tooth decay. *Pediatr Dent* 1997;19:22-7.
6. Huntington NL, Kim J, Hughes CV. Caries-risk factors for Hispanic children affected by early childhood caries. *Pediatr Dent* 2002;24:536-42.
7. Davies GN. Early childhood caries: A synopsis. *Community Dent Oral Epidemiol* 1998;26(suppl 1):106-16.
8. Berkowitz. Etiology of nursing caries: A microbiologic perspective. *J Public Health Dent* 1996;56:51-4.
9. Duchin S, Houte VJ. Relationship between *streptococcus mutans* and lactobacilli to incipient smooth surface dental caries in man. Oxford, UK: Pergamon Press Ltd; 1978:779-85.
10. Ramos-Gomez FJ, Weintraub JA, Gansky SA, Hoover CI, Featherstone JD. Bacterial, behavioral, and environmental factors associated with early childhood caries. *J Clin Pediatr Dent* 2002;26:165-73.
11. Shimono T, Sobue S. A new colorimetric caries activity test for diagnosis. *Dent Outlook* 1974;46:829-35.
12. Nishimura M, Bhuiyan MM, Matsumura S, Shimono T. Assessment of the caries activity test (Cariostat) based on the infection levels of mutans streptococci and lactobacilli in 2- to 13-year-old children's dental plaque. *J Dent Child* 1998;65:248-51.
13. Nishimura M, Docor OR, Chen HJ. The Cariostat as a screening method for approximal caries in primary dentition. *Pediatr Dent* 1993;3:35-49.
14. World Health Organization. Oral Health Surveys: Basic Methods. 4th ed. Geneva, Switzerland: WHO; 1997.
15. Milgrom P, Mancl L, King B, Weinstein P, Wells N, Jeffcott E. An explanatory model of dental care utilization of low-income children. *Med Care* 1998;36:554-66.
16. Wong MCM, Schwarz E, Lo ECM. Patterns of dental caries severity in Chinese kindergarten children. *Community Dent Oral Epidemiol* 1997;25:343-7.
17. Wong MCM, Lo ECM, Schwarz E, Zhang HG. Oral health status and oral health behaviors in Chinese children. *J Dent Res* 2001;80:1459-65.
18. Wennhall I, Mårtensson EM, Sjunnesson I, Matsson L, Schröder U, Twetman S. Caries-preventive effect of an oral health program for preschool children in a socioeconomic, multicultural area in Sweden: Results after one year. *Acta Odontol Scand* 2005;63:163-7.
19. Feldens CA, Vítolo MR, Drachler MDL. A randomized trial of the effectiveness of home visits in preventing early childhood caries. *Community Dent Oral Epidemiol* 2007;35:215-23.
20. Davies GM, Duxbury JT, Boothman NJ, Davies RM, Blinkhorn AS. A staged intervention dental health promotion program to reduce early childhood caries. *Community Dent Health* 2005;22:118-22.
21. Riedy CA, Weinstein P, Milgrom P. An ethnographic study for understanding children's oral health in a multicultural community. *Int Dent J* 2001;51:305-12.
22. Bu LJ, Xu Q, Zeng WQ. An analysis of etiology in primary tooth caries. *Chinese Primary Health Care* 2004;18:93-4.

23. Huang FX. The analysis of caries risk factors in 890 pre-school children. *Anthol Med* 2000;19:367-8.
24. Marino R, Bonze K, Scholl T, Anhalt H. Nursing bottle caries: Characteristics of children at risk. *Clin Pediatr* 1989;28:129-31.
25. American Academy of Pediatric Dentistry. Policy on early childhood caries (ECC): Classifications, consequences, and preventive strategies. Reference Manual 2003-04. *Pediatr Dent* 2003;25:24-6.
26. Reisine ST, Psoter W. Socioeconomic status and selected behavioral determinants as risk factors for dental caries. *J Dent Educ* 2001;65:1009-16.
27. King NM, Wu II, Tsai JS. Caries prevalence and distribution and oral health habits of 0- to 4-year-old children in Macau, China. *J Dent Child (Chic)* 2003;70:243-7.
28. Livny A, Sgan-Cohen HD. A review of a community program aimed at preventing early childhood caries among Jerusalem infants: A brief communication. *J Health Dent* 2007;67:78-84.
29. Jin BH, Ma DS, Moon HS, Paik DI, Hahn SH, Horowitz AM. Early childhood caries: Prevalence and risk factors in Seoul, Korea. *J Health Dent* 2003;63:183-8.
30. Acs G, Lodolini G, Kaminsky S, Cisneros GJ. Effect of nursing caries on body weight in a pediatric population. *Pediatr Dent* 1992;14:302-5.
31. Ayhan H, Suskan E, Yildirim S. The effect of nursing or rampant caries on height, body weight, and head circumference. *J Clin Pediatr Dent* 1996;20:209-12.
32. Featherstone J. The science and practice of caries prevention. *J Am Dent Assoc* 2000;131:887-99.
33. Winston AE. Caries prevention in the 21st century. *J Am Dent Assoc* 1998;129:1579-87.
34. Loesche WJ. Role of *Streptococcus mutans* in human dental decay. *Microbiol Rev* 1986;50:353-80.
35. Leverett DH, Proskin HM, Featherston JD, et al. Caries risk assessment in a longitudinal discrimination study. *J Dent Res* 1993;72:538-43.
36. Leverett DH, Featherstone JD, Proskin HM, et al. Caries risk assessment by a cross-sectional discrimination model. *J Dent Res* 1993;72:529-37.
37. Ansai T, Yamashita Y, Shibata Y, et al. Relationship between dental caries experience of a group of Japanese kindergarten children and the result of two caries activity tests conducted on their saliva and dental plaque. *Int J Paediatr Dent* 1994;4:13-7.

## Abstract of the Scientific Literature

### Psychological outcome of orthodontic treatment

*The purpose of this study was to investigate oral health, social, and psychological effects of malocclusion, and the effectiveness of orthodontic treatment. This study began in 1981 with 1018 participants ages 11-12 years. Assessments of dental health and psychosocial well-being were conducted based on facial and dental photographs, study casts, and questionnaires. The photos and casts were rated for attractiveness and treatment need. Questionnaires were given at the beginning of the study as well as at each subsequent visit, investigating the relationships between occlusal status, attractiveness, psychological well-being, social status and achievement, and quality of life. Individual interviews were also conducted. No recommendations regarding orthodontic treatment were made and the subjects were observed over time. The subjects either received orthodontic treatment or not based on their own volition. At the third follow-up in 2001, 337 subjects between the ages of 30-31 were reexamined and one-way ANOVA was used to explore difference between 4 groups: treatment need vs. no need, and treatment received vs. no treatment. Although participants with a prior need who obtained treatment had better tooth alignment and satisfaction, when self-esteem at baseline was controlled for, orthodontic treatment had little positive impact on psychological health and quality of life in adulthood.*

**Comments:** *This is the first in a series of articles from a 20-year cohort study of health gain from orthodontic treatment conducted in the United Kingdom. It is interesting to know that when pre-treatment self-esteem was accounted for, all the health gains in the psychological aspect vanished. It would seem that the potential impact of orthodontic treatment on psychological health has more to do with the patient's initial self-esteem rather than whether the patient received treatment. In the United States, where the youth might be more image-conscious, it is possible that a duplicate study will affirm the results expressed here. RHH*

*Address correspondence to Dr. W. C. Shaw, School of Dentistry, University of Manchester, Manchester M15 6FH, United Kingdom; e-mail: bill.shaw@manchester.ac.uk*

**Shaw WC, Richmond S, Kenealy PM, Kingdon A, Worthington H. A 20-year cohort study of health gain from orthodontic treatment: Psychological outcome. *Am J Orthod Dentofacial Orthop* 2007;132:146-57.**

49 references